LISTING OF THE CLAIMS

The following is a marked-up version of the Claims pursuant to 37 C.F.R. §1.121 (c)(1)(ii) with instructions and markings showing changes made herein to the previous version of the Claims on record. Underlining denotes added text while strikeout denotes deleted text.

- (Withdrawn) An enzymatically active variant of a precursor phenol oxidizing enzyme, said precursor being derived from a Stachybotrys species.
- 2. (Withdrawn) An enzymatically active variant of a precursor phenol oxidizing enzyme, which precursor phenol oxidizing enzyme has at least 68% identity to the amino acid sequence as disclosed in SEQ ID NO:2, said variant comprising a sequence that differs from that of said precursor in at least one of the positions 48, 67, 70, 76, 83, 98, 115, 119, 134, 171, 175, 177, 179, 188, 236, 246, 253, 254, 269, 272, 296, 302, 308, 318, 329, 331, 346, 348, 349, 365, 390, 391, 394, 404, 415, 423, 425, 428, 434, 465, 479, 481, 483, 499, 550, 562, 570, and 573, or sequence positions corresponding thereto.
- 3. (Withdrawn) The phenol oxidizing enzyme variant of claim 2, wherein said variant comprises a sequence that differs from that of said precursor in at least one of the positions 254, 272, 346, 348, 394, and 425, or sequence positions corresponding thereto.
- 4. (Withdrawn) The phenol oxidizing enzyme variant of claim 2 including an amino acid substitution at one or more of the positions or position sets: 76/254/302; 76/254/302/188; 76/254/302/394/425; 119/254/329; 119/254/390; 119/254/415; 171/254/346; 236/254; 254, 254/272; 254/302/346/348; 254/346/348; 254/394; 254/550; and 394/425.

(Withdrawn) The phenol oxidizing enzyme variant of claim 2 having at least one 5. amino acid substitution or substitution set selected from:

N391S
G115S
D562G
D394N/V425M
V134I/H1 <u>77Y</u>
L499F
M254F
M254F/L499F
M98L/M254F
L76W/M264F
M254F/F349Y
H175V
H177V
L76W/M254F/E302V
M254F/D394N/V425M
L76W/M254F/E302V/D394N/V425M
M254F/A296S
M254F/W318Y
M254F/L48Y
M254F/R83K
M254F/M188F
M254F/Q246H
M254F/\$331T
M254F/V483T
M254F/R67T
V119L/M254F/N70V
M254F/N70V
M254F/D308\$
M254F/E365T
M254F/\$415A
M254F/R423A
M254F/D428G
M254F/R434E
M254F/E465M
M254F/A <u>479G</u>
M254F/N550A
P253A
V119L/M254F/A269M
M254F/A269M
V119L/M254F/G329N
M254F/G329N
M254F/S331A

M254F/E346V/E348Q
V119L/M254F/E346V
M254F/E346V
V119L/M254F/A390P
M254F/A390P
M254F/N404T
V119L/M254F/\$415L
M254F/S415L
M254F/R481G
M254F/A573N
M254F/A573N/F570L
M254F/L76W/E302V/M188K
M254N
M254L
M254A
M2541
M254E
M254S
M254H
M254V
M254T
M254P
M254G
M254K
M254C
M254F/D394G
M254F/D394V
M254F/D394S
M254F/D394H
M254F/D394P
M254F/D394Y
M254F/D394W
M254F/D394N
M254F/M179F
M254F/M179V
M254F/M179P
M254F/M179G
M254F/M179E
M254F/M179L
M254F/1181D
M254F/\$180F/l181L
M254F/E346V/E302I
M254F/E346V/E302K
M254F/E346V/E348Q/E302F

M254F/E346V/E348Q/E302A	
M254F/E346V/E348Q/E302L	
M254F/E346V/E302C	
M254F/E346V/E302V	
M254F/E346V/M171T	_
M254F/E346V/E348Q/M171P	_
M254F/E346V/E348Q/M171L	_
M254F/E346V/M171Y	
M254F/ E346V/E348Q/M171V	_
M254F/ E346V/M171S	
M254F/E346V/M171R	
M254F/E346V/E348Q/M171F	
M254F/E346V/ M171K	
M254F/E346V/E348Q/M171Q	
M254F/E346V/E348Q/M171N	_
M254F/E346V/E348Q/M171N/L172H	
M254F/S272L	
M254F/E236K	
M254F/E346V/E348Q/M188K/D394W/\$272L/E23	3ř
M254F/E346V/E348Q/M188K/D394W/E236Q	_
M254F/E346V/E348Q/M188K/D394W/E236K	_
M254F/E346V/E348Q/M188K/D394W/E236D	
M254F/E346V/E348Q/M188K/D394W/E236A	
M188K/M254F/E346V/E348Q/D394W	_

- 6. (Withdrawn) The phenol oxidizing enzyme variant of any one of claims 2, 3, 4, or 5, wherein said precursor phenol oxidizing enzyme has at least 80% identity, and preferably at least 85% identity, to the amino acid sequence disclosed in SEQ ID NO:2.
- 7. (Withdrawn) The phenol oxidizing enzyme variant of any one of claims 2, 3, 4, or 5, wherein said precursor phenol oxidizing enzyme has at least 90% identity, and preferably at least 95% identity, to the amino acid sequence disclosed in SEQ ID NO:2.
- 8. (Withdrawn) The phenol oxidizing enzyme variant of claim 7, wherein said precursor phenol oxidizing enzyme has the amino acid sequence disclosed in SEQ ID NO:2.

(Withdrawn) The phenol oxidizing enzyme variant of claim 2, wherein said phenol oxidizing enzyme variant has increased phenol oxidizing activity at high pH as compared to said precursor phenol oxidizing enzyme.

- 10. (Withdrawn) The phenol oxidizing enzyme variant of claim 9, having a pH optimum of at least 8, and preferably at least 9.
- 11. (Withdrawn) The phenol oxidizing enzyme variant of claim 2, wherein said precursor is obtainable from a Stachybotrys species, preferably Stachybotrys chartarum.
- 12. (Currently Amended) An isolated polynucleotide encoding an enzymatically active variant of a phenol oxidizing enzyme, wherein said phenol oxidizing enzyme has [at least 68% identity to] the amino acid sequence as disclosed in SEQ ID NO:2, said variant comprising a sequence that differs from that of said phenol oxidizing enzyme in at least one of the positions 48, 67, 70, 76, 83, 98, 115, 119, 134, 171, 175, 177, 179, 188, 236, 246, 253, 254, 269, 272, 296, 302, 308, 318, 329, 331, 346, 348, 349, 365, 390, 391, 394, 404, 415, 423, 425, 428, 434, 465, 479, 481, 483, 499, 550, 562, 570, and 573, or sequence positions corresponding thereto.
- 13. (Original) An expression vector comprising the polynucleotide of claim 12.
- 14. (Original) A host cell comprising the expression vector of claim 13.
- 15. (Original) The host cell of claim 14, wherein said host cell is a filamentous fungus.
- 16. (Original) The host cell of claim 15, wherein said fungus is an Aspergillus species or a Trichoderma species.
- 17. (Withdrawn) An isolated protein having phenol oxidizing activity, which comprises an amino acid sequence having at least 68% and less than 100% identity to the amino acid sequence disclosed in SEQ ID NO:2, and which differs from said SEQ ID

NO:2 sequence in at least one of the positions 48, 67, 70, 76, 83, 98, 115, 119, 134, 171, 175, 177, 179, 188, 236, 246, 253, 254, 269, 272, 296, 302, 308, 318, 329, 331, 346, 348, 349, 365, 390, 391, 394, 404, 415, 423, 425, 428, 434, 465, 479, 481, 483, 499, 550, 562, 570, and 573, or sequence positions corresponding thereto.

18. (Currently Amended) A method for obtaining a phenol oxidizing enzyme variant from a Stachybotrys species, comprising the steps of:

mutagenizing a gene encoding a phenol oxidizing enzyme, which phenol oxidizing enzyme comprises an amino acid sequence having at least 68% identity to the amino acid sequence shown in SEQ ID NO:2;

introducing the mutant gene into a host strain whereby a transformed host strain is obtained;

growing said transformed host whereby said mutant gene is expressed and a phenol oxidizing enzyme variant, differing from said phenol oxidizing enzyme by one or more amino acid substitutions, is identified by recovering said variant and screening it for increased phenol oxidizing activity and/or increased pH optimum.

- 19. (Original) The method of claim 18, wherein said one or more amino acid substitutions correspond to amino acid positions selected from the group consisting of 48, 67, 70, 76, 83, 98, 115, 119, 134, 171, 175, 177, 179, 188, 236, 246, 253, 254, 269, 272, 296, 302, 308, 318, 329, 331, 346, 348, 349, 365, 390, 391, 394, 404, 415, 423, 425, 428, 434, 465, 479, 481, 483, 499, 550, 562, 570, and 573 of said SEQ ID NO:2 sequence.
- 20. (Currently Amended) The method of claim 18, wherein said mutagenized gene is a cloned Stachybotrys gene , preferably a cloned Stachybotrys charterum gene, or a cloned gene capable of hybridizing to such a Stachybotrys gene under conditions of intermediate to high-stringency.
- 21. (Currently Amended) A method for producing a variant of a phenol oxidizing enzyme, which enzyme comprises an amino acid sequence having at least 68% identity

te-the amino acid sequence shown in SEQ ID NO:2; said method comprising the steps of:

- a) culturing a host cell comprising a polynucleotide encoding said variant, wherein said variant differs from said phenol oxidizing enzyme sequence in at least one of the positions 48, 67, 70, 76, 83, 98, 115, 119, 134, 171, 175, 177, 179, 188, 236, 246, 253, 254, 269, 272, 296, 302, 308, 318, 329, 331, 346, 348, 349, 365, 390, 391, 394, 404, 415, 423, 425, 428, 434, 465, 479, 481, 483, 499, 550, 562, 570, and 573, or sequence positions corresponding thereto, under conditions suitable for the production of said variant; and
 - (b) optionally recovering said variant produced.
- 22. (Withdrawn) A method for producing a host cell comprising a polynucleotide encoding a variant of a precursor phenol oxidizing enzyme, which precursor enzyme comprises an amino acid sequence having at least 68% identity to the amino acid sequence shown in SEQ ID NO:2; said method comprising the steps of:
- (a) obtaining a polynucleotide encoding said variant, wherein said variant differs from said precursor sequence in at least one of the positions 48, 67, 70, 76, 83, 98, 115, 119, 134, 171, 175, 177, 179, 188, 236, 246, 253, 254, 269, 272, 296, 302, 308, 318, 329, 331, 346, 348, 349, 365, 390, 391, 394, 404, 415, 423, 425, 428, 434, 465, 479, 481, 483, 499, 550, 562, 570, and 573, or sequence positions corresponding thereto;
 - (b) introducing said polynucleotide into said host cell; and
- (c) growing said host cell under conditions suitable for the production of said variant.
- 23. (Withdrawn) The method of claim 22, wherein said host cell is a filamentous fungus.
- 24. (Withdrawn) The host cell of claim 23, wherein said fungus is a Aspergillus species or a Trichoderma species.

PØ14

- 25. (Previously presented) The isolated polynucleotide of claim 12, wherein said polynucleotide encodes a variant which comprises a sequence that differs from that of the phenol oxidizing enzyme in at least one of the positions 254, 272, 346, 348, 394, and 425, or sequence positions corresponding thereto.
- 26. (Previously presented) The isolated polynucleotide of claim 12 wherein said polynucleotide encodes an amino acid substitution at one or more of the positions or position sets: 76/254/302; 76/254/302/188; 76/254/302/394/425; 119/254/329; 119/254/390; 119/254/415; 171/254/346; 236/254; 254; 254/272; 254/302/346/348; 254/346/348; 254/394; 254/550; and 394/425.
- 27. (Previously presented) The isolated polynucleotide of claim 12 encoding a variant having at least one amino acid substitution or substitution set selected from:

N391S
G115S
D562G
D394N/V425M
V134I/H177Y
L499F
M254F
M254F/L499F
M98L/M254F
L76W/M254F
M254F/F3 <u>49Y</u>
H175V
H177V
L76W/M254F/E302V
M254F/D394N/V425M
L76W/M254F/E302V/D394N/V425M
M254F/A296S
M254F/W318Y
M254F/L48Y
M254F/R83K
M254F/M188F
M254F/Q246H
M254F/S331T
M254F/V483T
M254F/R67T
V119L/M254F/N70V

M254F/N70V
M254F/D308S
M254F/E365T
M254F/S415A
M254F/R423A
M254F/D428G
M254F/R434E
M254F/E465M
M254F/A479G
M254F/N550A
P253A
V119L/M254F/A269M
M254F/A269M
V119L/M254F/G329N
M254F/G329N
M254F/S331A
M254F/E346V/E348Q
V119L/M254F/E346V
M254F/E346V
V119L/M254F/A390P
M254F/A390P
M254F/N404T
V119L/M254F/S415L
M254F/S415L
M254F/R481G
M254F/A573N
M254F/A573N/F570L
M254F/L76W/E302V/M188K
M254N
M254L
M254A
M254I
M254E
M254S
M254H
M254V
M254T
M254P
M254G
M254K
M254C
M254F/D394G
M254F/D394V
M254F/D394S
MZ04F/D3945

Ø016

M254F/D394H
M254F/D394P
M254F/D394Y
M254F/D394W
M254F/D394N
M254F/M179F
M254F/M179V
M254F/M179P
M254F/M179G
M254F/M179E
M254F/M179L
M254F/I181D
M254F/S180F/I181L
M254F/E346V/E302I
M254F/E346V/E302K
M254F/E346V/E348Q/E302F
M254F/E346V/E348Q/E302A
M254F/E346V/E348Q/E302L
M254F/E346V/E302C
M254F/E346V/E302V
M254F/E346V/M171T
M254F/E346V/E348Q/M171P
M254F/E346V/E348Q/M171L
M254F/E346V/M171Y
M254F/ E346V/E348Q/M171V
M254F/ E346V/M171S
M254F/E346V/M171R
M254F/E346V/E348Q/M171F
M254F/E346V/ M171K
M254F/E346V/E348Q/M171Q
M254F/E346V/E348Q/M171N
M254F/E346V/E348Q/M171N/L172H
M254F/S272L
M254F/E236K
M254F/E346V/E348Q/M188K/D394W/S272L/E236K
M254F/E346V/E348Q/M188K/D394W/E236Q
M254F/E346V/E348Q/M188K/D394W/E236K
M254F/E346V/E348Q/M188K/D394W/E236D
M254F/E346V/E348Q/M188K/D394W/E236A
M188K/M254F/E346V/E348Q/D394W

Please cancel claim 28.

Please cancel claim 29.

- 30. (Currently Amended) The isolated polynucleotide of claim 29 12, 25, 26 or 27, wherein the phenol oxidizing enzyme has the amino acid sequence disclosed in SEQ ID NO:2.
- 31. (Previously presented) The isolated polynucleotide of claim 12, wherein the phenol oxidizing enzyme variant has increased phenol oxidizing activity at high pH.
- 32. (Currently Amended) The isolated polynucleotide of claim 31, wherein the variant has a pH optimum of at least 8, and preferably at least 9.
- 33. (Currently Amended) The isolated polynucleotide of claim 12, wherein said phenol oxidizing enzyme is obtainable from a Stachybotrys species, preferably Stachybotrys chartarum.

Please add the following new claims.

- 34. (New) The method according to claim 20, wherein said cloned Stachybotrys chartarum gene is a cloned Stachybotrys chartarum gene or a cloned gene capable of hybridizing to such a Stachybotrys gene under conditions of intermediate to high stringency.
- 35. (New) The method according to claim 32, wherein the variant has a pH of at least 9.
- 36. (New) The method according to claim 33, wherein the Stachybotrys species is Stachybotrys chartarum.

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not infinted to the items enceked.
□ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
□ other.

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.